

ONE PACKS

Dispenser-Free Control Systems®

Sanitergent™ Cleaner/Sanitizer

#2507643

Specialized Sanitizer and Cleaner for:
Milkshake and Soft Serve Machines, Tanks,
Vats, Food Processing Equipment, Dishes,
Glasses, Utensils, Tabletops, Floors, and Walls

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF SWALLOWED: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED: Move person to fresh air. If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

DIRECTIONS



It may be necessary to remove gross filth and heavy soil from surface by a pre-scrub, pre-flush and where necessary, a pre-soak treatment. Dissolve 0.10 oz. of Sanitergent™ Cleaner/Sanitizer per 32 ounces of water to give a solution containing 100 ppm available chlorine. Apply the sanitizing solution to the surface to be sanitized spray to wet all surfaces thoroughly. Let stand at least one minute. Air dry. The sanitizing solution should be used promptly after mixing. It should be prepared daily or more often if it becomes dilute or soiled. If the solution is allowed to stand and/or continued use shows the level of chlorine has not been maintained, add more Sanitergent™ Cleaner/Sanitizer to maintain a use solution of 100 ppm available chlorine. Empty and rinse bottle before refilling.

EPA Reg. No. 10634-27-3640
EPA Est. No. 3640-WI-1

SANITERGENT™ CLEANER/SANITIZER
ST-764 FOR MILKSHAKE MACHINES, SOFTSERVE MACHINES AND FOOD PROCESSING UTENSILS
EPA Reg. No. 10634-27-3640 EPA Est. No. 3640-WI-1

ACTIVE INGREDIENT
Sodium dichloro-s-triazetrone dihydrate... 9%
NETT INGREDIENTS... 99%
TOTAL... 100%

Provides 3.3% available chlorine

DIRECTIONS FOR USE:
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Dissolve 0.10 oz. Sanitergent Cleaner/Sanitizer per 32 oz. of water to give a solution containing 100 ppm available chlorine. Please see outer carton for detailed directions.

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO
Highly Corrosive. Causes irreversible eye damage. Causes burns to skin, irritating to nose and throat. May be fatal if swallowed.
See outer carton for additional precautionary and first aid statements.

24-HOUR EMERGENCY CONTACT: 1-800-235-3924
SOLD BY: STEARNS PACKAGING CORPORATION,
4200 Sycamore Ave., Madison, WI 53714
Net Weight: 0.10 wt. oz. (2.83 g)

ONE PACKS

Dispenser-Free Control Systems®

Sanitergent™ Cleaner/Sanitizer


#2507643

Desinfectante y Limpiador Especializado para:
Batido de Leche y Máquinas de Servicio Blando,
Tanques, Cubas, Equipo de Procesamiento de
Alimentos, Platos, Vasos, Utensilios, Tableros
de Mesa, Pisos y Paredes

PRIMEROS AUXILIOS

SI ESTÁ EN LOS OJOS: Mantenga abierto el ojo y enjuague lenta y suavemente con agua durante 15-20 minutos. Retire los lentes de contacto, si los tiene, después de los primeros 5 minutos. Luego continúe enjuagando el ojo.
EN CASO DE CONTACTO CON LA PIEL O LA ROPA: Quite la ropa contaminada. Enjuague la piel inmediatamente con abundante agua durante 15-20 minutos. Llame a un centro de control de envenenamiento o a un médico de inmediato para recibir asesoramiento sobre el tratamiento.
EN CASO DE INGESTIÓN: Llame a un centro de control de envenenamiento o a un médico de inmediato para recibir asesoramiento sobre el tratamiento. Haga que la persona beba un vaso de agua si puede tragar. No induzca el vómito a menos que así lo indique un centro de control de envenenamientos o un médico. No le dé nada por la boca a una persona inconsciente.
EN CASO DE INHALACIÓN: Traslade a la persona al aire libre. Si la persona no respira, llame al 911 o a una ambulancia, luego dele respiración artificial, preferiblemente de boca a boca, si es posible. Llame a un centro de control de envenenamiento o a un médico de inmediato para recibir asesoramiento sobre el tratamiento.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.
NOTA PARA EL MÉDICO: El daño probable de la mucosa puede contraindicar el uso del lavado gástrico.

INTRUCCIONES



Puede ser necesario eliminar la suciedad gruesa y el suelo pesado de la superficie mediante un predesprendimiento, prelavado y, cuando sea necesario, un tratamiento de preempapado. Disuelva 0.10 oz de Sanitergent™ Cleaner / Sanitizer por 32 onzas de agua para dar una solución que contiene 100 ppm de cloro disponible. Aplique la solución desinfectante a la superficie para rociar desinfectado para humedecer completamente todas las superficies. Deje reposar al menos un minuto. Secar al aire. La solución desinfectante se debe usar inmediatamente después de mezclar. Debe prepararse diariamente o con mayor frecuencia si se diluye o se ensucia. Si la solución se deja en reposo y / o el uso continuado muestra que el nivel de cloro no se ha mantenido, agregue más Sanitergent™ Cleaner / Sanitizer para mantener una solución de uso de 100 ppm de cloro disponible. Vacíe y enjuague la botella antes de volver a llenarla.



STEARNS PACKAGING CORPORATION
P.O. Box 14495, 4200 Sycamore Avenue, Madison, Wisconsin 53708
1-800-655-5008 • www.stearnspkg.com

Rev. 0221



STEARNS PACKAGING CORPORATION
P.O. Box 14495, 4200 Sycamore Avenue, Madison, Wisconsin 53708
1-800-655-5008 • www.stearnspkg.com

Rev. 0221

EPA Reg. No. 10634-27-3640
EPA Est. No. 3640-WI-1

SANITERGENT™ CLEANER/SANITIZER
ST-764 FOR MILKSHAKE MACHINES, SOFTSERVE MACHINES AND FOOD PROCESSING UTENSILS
EPA Reg. No. 10634-27-3640 EPA Est. No. 3640-WI-1

ACTIVE INGREDIENT
Sodium dichloro-s-triazetrone dihydrate... 9%
NETT INGREDIENTS... 99%
TOTAL... 100%

Provides 3.3% available chlorine

DIRECTIONS FOR USE:
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Dissolve 0.10 oz. Sanitergent Cleaner/Sanitizer per 32 oz. of water to give a solution containing 100 ppm available chlorine. Please see outer carton for detailed directions.

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO
Highly Corrosive. Causes irreversible eye damage. Causes burns to skin, irritating to nose and throat. May be fatal if swallowed.
See outer carton for additional precautionary and first aid statements.

24-HOUR EMERGENCY CONTACT: 1-800-235-3924
SOLD BY: STEARNS PACKAGING CORPORATION,
4200 Sycamore Ave., Madison, WI 53714
Net Weight: 0.10 wt. oz. (2.83 g)